

LabDiet® JL Rat and Mouse/Auto 6F 5K52* 5K67*

DESCRIPTION

LabDiet® JL Rat and Mouse/Auto 6F diets are controlled under the Constant Nutrition® formula management program. 5K52 is the breeding diet used at The Jackson Laboratory. 5K67 is the same formula as 5K52, but in a standard oval shape.

Features and Benefits

- 5K52 is the primary breeding diet used at The Jackson Laboratory. Specific information on strains fed can be obtained from The Jackson Laboratory.
- 5K67, in addition to irradiated versions, is available to help your animals maintain consistency from the breeding colony.
- Designed to enhance breeding efficiency.
- Fortified with extra nutrients to compensate for losses during autoclaving
- Processed with silicon dioxide to reduce sticking and clumping

Product Forms Available

- 5K52: Cylinder shaped pellet - 3/8" diameter by 3/4" length
- 5K67: Oval shaped - 3/8" x 5/8" x 1" length

Other Versions Available

- 5LG4/5LL4 JL Rat and Mouse Irr 6F

GUARANTEED ANALYSIS

Crude protein not less than	18.0%
Crude fat not less than	6.0%
Crude fiber not more than	5.0%
Ash not more than	8.0%

INGREDIENTS

Ground wheat, ground corn, wheat middlings, ground oats, fish meal, dehulled soybean meal, soybean oil, corn gluten meal, dehydrated alfalfa meal, dicalcium phosphate, monocalcium phosphate, brewers dried yeast, calcium carbonate, menadione dimethylpyrimidinol bisulfite, salt, DL-methionine, choline chloride, magnesium oxide, thiamin mononitrate, pyridoxine hydrochloride, cholecalciferol, vitamin A acetate, calcium pantothenate, ferrous sulfate, biotin, manganous oxide, dl-alpha tocopheryl acetate, folic acid, vitamin B₁₂ supplement, riboflavin, nicotinic acid, zinc oxide, ferrous carbonate, copper sulfate, zinc sulfate, cobalt carbonate, calcium iodate.

FEEDING DIRECTIONS

Feed ad libitum to rodents. Provide plenty of fresh clean water at all times.

AUTOCLAVING SUGGESTIONS

During the autoclaving process, the pellets can be placed on trays, in small bags or in larger bags, as long as the pellets are stacked no more than 3 inches high. When steam autoclaved, the pellets swell and exert force on adjacent pellets. If confined by a bag or container, the pressure causes sticking as greater polymerization of fibrous materials occurs under such conditions. **Assay before and after autoclaving:** Conditions of sterilization must be determined for each autoclaving unit. It is best to assay the diet before and after sterilization to determine nutrient losses. Microbiological studies should be done also to insure the degree of sterilization desired.

For Product Availability, visit www.labdiet.com.

CHEMICAL COMPOSITION

Nutrients**

Protein, %	19.3
Arginine, %	1.03
Cystine, %	0.25
Glycine, %	0.94
Histidine, %	0.44
Isoleucine, %	0.87
Leucine, %	1.52
Lysine, %	0.97
Methionine, %	0.73
Phenylalanine, %	0.85
Tyrosine, %	0.56
Threonine, %	0.68
Tryptophan, %	0.23
Valine, %	0.90
Serine, %	0.98
Aspartic Acid, %	1.87
Glutamic Acid, %	4.52
Alanine, %	1.13
Proline, %	1.53
Taurine, %	0.03
Fat (ether extract), %	6.2
Fat (acid hydrolysis), %	7.2
Cholesterol, ppm	240
Linoleic Acid, %	2.88
Linolenic Acid, %	0.37
Arachidonic Acid, %	0.01
Omega-3 Fatty Acids, %	0.46
Total Saturated Fatty Acids, %	1.24
Total Monosaturated Fatty Acids, %	1.37
Fiber (Crude), %	4.3
Neutral Detergent Fiber ³ , %	15.1
Acid Detergent Fiber ⁴ , %	5.2
Nitrogen-Free Extract (by difference), %	53.6
Starch, %	38.9
Glucose, %	0.12
Fructose, %	0.15
Sucrose, %	0.62
Lactose, %	0.00
Total Digestible Nutrients, %	76.3
Gross Energy, kcal/gm	4.17
Physiological Fuel Value⁵, kcal/gm	3.47
Metabolizable Energy, kcal/gm	3.17
Minerals	
Ash, %	6.5
Calcium, %	1.17
Phosphorus, %	0.93
Phosphorus (non-phytate), %	0.68
Potassium, %	0.66
Magnesium, %	0.22

Sulfur, %	0.33
Sodium, %	0.26
Chlorine, %	0.45
Fluorine, ppm	37
Iron, ppm	380
Zinc, ppm	85
Manganese, ppm	160
Copper, ppm	11
Cobalt, ppm	0.80
Iodine, ppm	2.1
Chromium, ppm	2.0
Selenium, ppm	0.30

Vitamins

Carotene, ppm	1.5
Vitamin K (as menadione), ppm	20
Thiamin Hydrochloride, ppm	79
Riboflavin, ppm	9.0
Niacin, ppm	90
Pantothenic Acid, ppm	37
Choline Chloride, ppm	2000
Folic Acid, ppm	1.9
Pyridoxine, ppm	10
Biotin, ppm	0.30
B ₁₂ , mcg/kg	50
Vitamin A, IU/gm	20
Vitamin D ₃ (added), IU/gm	4.3
Vitamin E, IU/kg	45
Ascorbic Acid, mg/gm	—

Calories provided by:

Protein, %	22.238
Fat (ether extract), %	16.028
Carbohydrates, %	61.734

*Product Code

1. Based on the latest ingredient analysis information. Since nutrient composition of natural ingredients varies, analysis will differ accordingly.
2. Nutrients expressed as percent of ration except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.
3. NDF = approximately cellulose, hemi-cellulose and lignin.
4. ADF = approximately cellulose and lignin.
5. Physiological Fuel Value (kcal/gm) = Sum of decimal fractions of protein, fat and carbohydrate (use Nitrogen Free Extract) x 4,9,4 kcal/gm respectively.